



For adults with Familial Chylomicronemia Syndrome (FCS),

# ALL THE PLACES YOU WILL LO

Get to know Redemplo® and  
if it's right for you

Not actual patients.

## INDICATION

REDEMPLO® (plozasiran) is an injectable prescription medicine used together with a low-fat diet to reduce triglycerides (fat in the blood) in adults with a condition that keeps the body from breaking down fats called familial chylomicronemia syndrome (FCS).

Please see additional [Important Safety Information](#) and [Important Product Information](#) and [Patient Product Information](#) for REDEMPLO.



Not an actual patient.

## What is FCS?

FCS is an extreme type of severe hypertriglyceridemia (sHTG) that makes it hard for your body to break down and remove triglycerides.

People with FCS may have **10 to 100 times more triglycerides** than normal in their blood, even with standard treatment and a strict low-fat diet (less than 20 grams of fat per day).

For full definitions, see the [FCS Glossary on page 14](#).




FCS can cause symptoms like **acute pancreatitis, abdominal pain, brain fog, eruptive xanthomas, and nausea**

## Recognizing the potential burden of FCS

Did you know that acute pancreatitis caused by high triglycerides can be more serious than other causes of acute pancreatitis? Even one event can increase the chance of another.

**Triglycerides over 500 mg/dL put people with FCS in an acute pancreatitis risk danger zone**



Retrospective cohort study of annualized incidence rate of acute pancreatitis. Data were obtained from IQVIA's US Ambulatory Electronic Medical Records database (N=7,119,195).

When triglycerides are **above 880 mg/dL**, the risk of acute pancreatitis can be about **14 times higher** than triglycerides under 200 mg/dL.

With FCS, **just one meal can spike your triglycerides** to this level, drastically increasing your risk.

To lower the risk, it's recommended to lower triglycerides to **below 500 mg/dL**.



## Redemplo<sup>®</sup> is a treatment for adults with FCS that works in combination with diet to reduce triglycerides

A new **LO** starts now—data showed patients taking Redemplo saw a significant reduction in triglycerides, and a lowered number of acute pancreatitis attacks, with just one dose every 3 months.



### 80% Reduction in Triglycerides\*

A significant median<sup>†</sup> reduction of **80%** in triglycerides from baseline was seen at 10 months



### Lowered Acute Pancreatitis Attacks<sup>‡</sup>

With Redemplo, fewer people had acute pancreatitis attacks: 2 (8%) compared with 5 (20%) on placebo

\*Redemplo 25 mg lowered typical triglyceride levels by 80% (to 443 mg/dL) compared with 17% for placebo (to 1614 mg/dL).

<sup>†</sup> The median is the middle result in a group of people. This means that half of the patients had results lower than this number and half had results that were higher.

<sup>‡</sup> The clinical trial included an evaluation of acute pancreatitis events that occurred in patients taking Redemplo compared to those taking placebo. Redemplo has not been proven to reduce acute pancreatitis events.


The safety and effectiveness of Redemplo were studied in a **randomized, double-blind, placebo-controlled clinical trial** in 75 adult patients with FCS in combination with a low-fat diet (less than or equal to 20 grams of fat per day). The main goal of the trial was to examine how much Redemplo lowered triglycerides from the start of the trial to Month 10 compared with placebo.

Please see additional [Important Safety Information](#) and [Important Product Information](#) and [Patient Product Information](#) for REDEMPLO.



## Redemplo<sup>®</sup> helps lower triglyceride levels and keep them down

### Significant and Sustained Triglyceride Reduction With Redemplo 25 mg\*



## Most people in the study were able to stay on Redemplo

In the clinical study, the most common side effects were an increase in blood sugar (hyperglycemia), headache, nausea, and injection site reactions.


### IMPORTANT SAFETY INFORMATION

**Before you start using REDEMPLO**, tell your healthcare provider about all your medical conditions, including if you are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. It is not known if REDEMPLO could harm your unborn baby, or if it passes into your breast milk and could harm your breastfeeding baby.


Tell your healthcare provider or pharmacist about any prescription and over-the-counter medicines, vitamins, or herbal supplements you take. (Cont'd on page 7)

## How Redemplo<sup>®</sup> can help you reach a new **LO**


**When taken in addition to a low-fat diet, Redemplo helps reduce fat buildup in the blood and lowers high triglycerides**



People with FCS have very high triglycerides (fat in the blood). This causes fat particles called chylomicrons to build up in the bloodstream, which can make the blood look milky or cloudy.



APOC3 is a protein that sticks to triglyceride-rich chylomicrons in the blood. Sometimes it helps to think of APOC3 like a “speed bump” that slows down how quickly the body can clear these fat particles.



Redemplo uses a small piece of RNA, called siRNA, to reduce the amount of APOC3 your liver makes. With less APOC3, your body can clear fat from your blood more effectively.

**For full definitions, see the FCS Glossary on page 14**

APOC3, apolipoprotein C-III; RNA, ribonucleic acid; siRNA, small interfering RNA.

### IMPORTANT SAFETY INFORMATION (cont'd)

#### WHAT ARE THE POSSIBLE SIDE EFFECTS OF REDEMPLO?

The most common side effects of REDEMPLO include increased blood sugar levels, headache, nausea, and injection site reactions (pain, redness, or swelling).

These are not all the possible side effects of REDEMPLO. Tell your healthcare provider or treatment team if you have any side effect that bothers you or that does not go away.



**Redemplo<sup>®</sup> is just one convenient dose every 3 months, allowing for lasting results**



**When it comes to taking Redemplo, think 3:**

3 steps every 3 months



**Prepare**



**Inject**



**Dispose**

**Visit [Redemplo.com/how-to-inject](https://www.Redemplo.com/how-to-inject) to watch a full step-by-step video on how to take Redemplo.**

#### IMPORTANT SAFETY INFORMATION

**Before you start using REDEMPLO**, tell your healthcare provider about all your medical conditions, including if you are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. It is not known if REDEMPLO could harm your unborn baby, or if it passes into your breast milk and could harm your breastfeeding baby.

Tell your healthcare provider or pharmacist about any prescription and over-the-counter medicines, vitamins, or herbal supplements you take.



**When taking Redemplo<sup>®</sup>, it's important to stay on a low-fat diet (less than or equal to 20 grams of fat per day).**

Please see additional [Important Safety Information](#) and [Important Product Information](#) and [Patient Product Information](#) for REDEMPLO. 9



Not an actual patient.



## Support and resources are available with the Rely On Redemplo<sup>®</sup> Patient Support Program

The Rely On Redemplo Patient Support Program provides support, education, and information on financial assistance options that may be available to help you start and stay on track with Redemplo<sup>®</sup> (plozasiran).

**If you have been prescribed Redemplo<sup>®</sup>, you may pay as little as \$0 out of pocket (OOP) for Redemplo 25 mg with the Rely On Redemplo Copay Card\***

Eligible commercially insured patients may qualify to lower their OOP costs for Redemplo.

\*Eligibility requirements and terms and conditions apply.

### Enroll in the Rely On Redemplo Patient Support Program!

Your doctor can prescribe Redemplo and start your enrollment by completing the Redemplo Start Form for you. Then you'll be able to provide your consent for enrollment by signing digitally or in person.

**For support, call 1-844-REDEMPLO (1-844-733-3675), Monday to Friday, 8 AM to 8 PM ET.**

Please see additional [Important Safety Information](#) and [Important Product Information and Patient Product Information for REDEMPLO](#).





## FCS Glossary

Here are a few key terms to help you better understand FCS and how treatment like Redemplo<sup>®</sup> works.

**Acute pancreatitis:** Inflammation of the pancreas that appears suddenly.

**APOC3 (apolipoprotein C-III):** A protein the liver creates which slows the breakdown and removal of triglyceride-rich particles (fat particles) from the bloodstream. Think of APOC3 as a “speed bump” that slows down how quickly the body can clear these fat particles.

**Chylomicrons:** Large triglyceride-rich particles made in the intestines that carry fat from the food you eat through the bloodstream.

**Eruptive xanthomas:** Firm, raised, fatty lumps on the body.

**FCS:** An extreme type of severe hypertriglyceridemia that makes it hard for your body to break down and remove triglycerides. It may lead to serious health problems, such as sudden and painful acute pancreatitis events, fatigue and physical weakness, and eruptive xanthomas.

**Hyperglycemia:** An increase in blood sugar levels.

**Incidence:** The occurrence or frequency at which something occurs.

**Median:** The middle result in a group of people. This means that half of the patients had results lower than this number and half had results that were higher.

**mRNA (messenger RNA):** mRNA carries the instructions from DNA that tell the cell which protein to make.

**Placebo-controlled trial:** A clinical study in which some participants receive the study medicine (in this case, Redemplo) and others receive an inactive substance (called a placebo) that looks the same but has no effect. This allows researchers to understand whether changes are truly from the study medicine.

**siRNA (small interfering RNA) therapy:** Targets and destroys specific mRNA molecules, stopping the cell from making a certain protein.

### INDICATION

REDEMPLO<sup>®</sup> (plozasiran) is an injectable prescription medicine used together with a low-fat diet to reduce triglycerides (fat in the blood) in adults with a condition that keeps the body from breaking down fats called familial chylomicronemia syndrome (FCS).

### IMPORTANT SAFETY INFORMATION

**Before you start using REDEMPLO,** tell your healthcare provider about all your medical conditions, including if you are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. It is not known if REDEMPLO could harm your unborn baby, or if it passes into your breast milk and could harm your breastfeeding baby.

Tell your healthcare provider or pharmacist about any prescription and over-the-counter medicines, vitamins, or herbal supplements you take.

### WHAT ARE THE POSSIBLE SIDE EFFECTS OF REDEMPLO?

The most common side effects of REDEMPLO include increased blood sugar levels, headache, nausea, and injection site reactions (pain, redness, or swelling).

These are not all the possible side effects of REDEMPLO. Tell your healthcare provider or treatment team if you have any side effect that bothers you or that does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/safety/medwatch](http://www.fda.gov/safety/medwatch), or call 1-800-FDA-1088.

**Ask your doctor if Redemplo<sup>®</sup> is right for you**

Sign up for the latest updates and resources from Redemplo.

Visit [Redemplo.com/sign-up](http://Redemplo.com/sign-up)



## Your new **LO** starts with Redemplo<sup>®</sup>, with just one dose every 3 months



### 80% Reduction in Triglycerides\*

A significant median<sup>†</sup> reduction of **80%** in triglycerides from baseline was seen at 10 months



### Lowered Acute Pancreatitis Attacks<sup>‡</sup>

With Redemplo, fewer people had acute pancreatitis attacks: 2 (8%) compared with 5 (20%) on placebo



### Convenient Dosing Schedule

Redemplo is the only treatment that is just one dose every 3 months

The safety and effectiveness of Redemplo were studied in a randomized, double-blind, placebo-controlled clinical trial in 75 adult patients with FCS in combination with a low-fat diet (less than or equal to 20 grams of fat per day).

\*Redemplo 25 mg reduced median triglycerides by 80% (to 443 mg/dL) compared with 17% for placebo (to 1614 mg/dL).

<sup>†</sup>The median is the middle result in a group of people. This means that half of the patients had results lower than this number and half had results that were higher.

<sup>‡</sup>The clinical trial included an evaluation of acute pancreatitis events that occurred in patients taking Redemplo compared to those taking placebo. Redemplo has not been proven to reduce acute pancreatitis events.

## IMPORTANT SAFETY INFORMATION

### WHAT ARE THE POSSIBLE SIDE EFFECTS OF REDEMPLO?

The most common side effects of REDEMPLO include increased blood sugar levels, headache, nausea, and injection site reactions (pain, redness, or swelling).

These are not all the possible side effects of REDEMPLO. Tell your healthcare provider or treatment team if you have any side effect that bothers you or that does not go away.

Please see additional [Important Safety Information](#) and [Important Product Information](#) and [Patient Product Information](#) for REDEMPLO.

Ask your doctor if **Redemplo** is right for you

Sign up for updates and to learn more at [Redemplo.com](https://www.Redemplo.com)